

Finitistic Dimension Conjecture and Conditions on Ideals

Jiaqun WEI *

Department of Mathematics, Nanjing Normal University

Nanjing 210097, P.R.China

Email: weijiaqun@njnu.edu.cn

Abstract

The notion of Igusa-Todorov classes is introduced in connection with the finitistic dimension conjecture. As application we consider conditions on special ideals which imply the Igusa-Todorov and other finiteness conditions on modules proving the finitistic dimension conjecture and related conjectures in those cases.

MSC: 16E05; 16E10; 16G20

Key words: finitistic dimension; Igusa-Todorov class; ideal; radical

Introduction

Throughout the paper, we work on artin algebras and finitely generated left modules.

Let A be an artin algebra. Recall that the finitistic dimension of A , denoted by $\text{findim} A$, is defined to be the supremum of the projective dimensions of all finitely generated modules of finite projective dimension. The famous finitistic dimension conjecture claims that $\text{findim} A < \infty$, for any artin algebra A . The conjecture is still open and is related to many other homological conjectures (e.g., the Gorenstein symmetry conjecture, the Wakamatsu-tilting conjecture and the generalized Nakayama conjecture).

Until recently, only a few classes of algebras were known to have finite finitistic dimension, see for instance [1, 2, 3] etc.. In particular, the finitistic dimension conjecture holds for algebras of representation dimension at most 3, algebras with radical cube zero, monomial algebras and left serial algebras, etc..

All the above-mentioned algebras are Igusa-Todorov algebras, a notion introduced in [12] for algebras whose finiteness of finitistic dimensions is assured by the Igusa-Todorov functor [5]. Many algebras which are recently proved to have finite finitistic dimension are also Igusa-Todorov [12].

In this paper, we introduce the notion of Igusa-Todorov classes and apply them to obtain new Igusa-Todorov algebras. The main results are as follows.

*Supported by the National Science Foundation of China (No.10601024)

Theorem 0.1 *Let A be an artin algebra.*

(1) *Assume that I, J are two ideals of A such that $I\text{Jrad}A = 0$. Then A is Igusa-Todorov provided one of the following conditions is satisfied.*

- (i) *A/I is representation-finite, $\text{pd}_A J < \infty$ and A/J is syzygy-finite.*
- (ii) *A/J is representation-finite, $\text{pd}_A I < \infty$ and A/I is syzygy-finite.*
- (iii) *Both A/I and A/J are syzygy-finite and both ${}_A I$ and ${}_A J$ have finite projective dimension.*

(2) *Assume that $0 = I_0 \subseteq I_1 \subseteq \cdots \subseteq I_n$ are ideals in A such that $\text{pd}_A I_{i+1} < \infty$ and $(I_{i+1}/I_i)\text{rad}(A/I_i) = 0$, for each $0 \leq i \leq n-1$. If A/I_n is syzygy-finite (resp., syzygy-bounded, Igusa-Todorov), then A is also syzygy-finite (resp., syzygy-bounded, Igusa-Todorov).*

In particular, the finitistic dimension of A is finite in all the above cases.

Recall now the following well-known conjectures (see for instance [3, 7]). Assume that A is an artin algebra.

Gorenstein symmetry conjecture $\text{id}({}_A A) < \infty$ if and only if $\text{id}(A_A) < \infty$, where id denotes the injective dimension.

Wakamatsu-tilting conjecture Let ${}_A \omega$ be a Wakamatsu-tilting module. (1) If $\text{pd}_A \omega < \infty$, then ω is tilting. (2) If $\text{id}_A \omega < \infty$, then ω is cotilting.

Generalized Nakayama conjecture Each indecomposable injective A -module occurs as a direct summand in the minimal injective resolution of ${}_A A$.

Note that the Gorenstein symmetry conjecture and the generalized Nakayama conjecture are special cases of the second Wakamatsu-tilting conjecture. Moreover, if the finitistic dimension conjecture holds for A and A° (hereafter A° denotes the opposite algebra of A), then all the above conjectures hold.

Small's results [10] shows that if A has a nilpotent ideal I such that $\text{pd}_A I < \infty$ and $(A/I)^\circ$ has finite finitistic dimension, then A° has also finite finitistic dimension. Hence, combining with the fact that, for a syzygy-finite algebra A , both the finitistic dimensions of A and A° are finite (see for instance [11]), we have the following corollary.

Corollary 0.2 *The Gorenstein symmetry conjecture, the Wakamatsu-tilting conjecture and the generalized Nakayama conjecture hold for algebras in Theorem 0.1 (1), provided that moreover, one of there considered ideals is nilpotent.*

In particular, Small's results [10] shows that the finitistic dimension of A° is finite provided that $\text{pd}_A \text{rad}^2 A < \infty$ or $\text{pd}_A \text{rad}^3 A < \infty$ (see for instance [19]). It is natural to ask if the finitistic dimension of A is also finite in these cases. The following corollary gives a partial answer to that question.

Corollary 0.3 *Let A be an artin algebra. Then the finitistic dimension of A is finite provided that A satisfies one of the following two conditions.*

- (1) *$\text{rad}^5 A = 0$ and $\text{pd}_A \text{rad}^2 A < \infty$.*
- (2) *$\text{rad}^n A = 0$ and $\text{pd}_A \text{rad}^i A < \infty$ for each $i \geq 3$.*

In particular, the Gorenstein symmetry conjecture, the Wakamatsu-tilting conjecture and the generalized Nakayama conjecture hold for algebras in two cases.

In the last section, we give examples to show that our results do apply to obtain algebras with finite finitistic dimension. In particular, our methods are compared with the others, for instance, the recent results of Huards, Lanzilotta and Mendoza [4] using their infinite radical.

1 Preliminaries

Throughout this paper, A is always an artin algebra. We denote by $A\text{-mod}$ the category of all (finitely generated) A -modules. If M is an A -module, we use $\text{pd}_A M$ to denote the projective dimension of ${}_A M$ and use $\Omega_A^i M$ to denote the i -th syzygy of M (we set $\Omega_A^0 M := M$). The notion $\text{add}_A M$ denotes the category consists of all direct summands of finite direct sums of M .

Let $\mathcal{C}$ be a class of A -modules. We denote $\text{gd}\mathcal{C} := \sup\{\text{pd}_A M \mid M \in \mathcal{C}\}$. We use $\Omega^n \mathcal{C}$ to denote the class of A -modules which are isomorphic to $\Omega_A^n M$ for some $M \in \mathcal{C}$. The class $\mathcal{C}$ is said to be *add-finite* provided that $\mathcal{C} \subseteq \text{add}_A C$ for some A -module C , or equivalently, the number of non-isomorphic indecomposable direct summands of objects in $\mathcal{C}$ is finite. Moreover, we say that $\mathcal{C}$ is *(n-)syzygy-finite* provided that there is some n such that $\Omega^n \mathcal{C}$ is add-finite. If $A\text{-mod}$ is syzygy-finite, then we also say that A is syzygy-finite. Note that if $\mathcal{C}$ is n -syzygy-finite, then $\Omega^i \mathcal{C}$ is add-finite for any $i \geq n$ and $\Omega^i \mathcal{C}$ is $(n-i)$ -syzygy-finite for each $1 \leq i < n$.

More generally, we say that a class of A -modules $\mathcal{C}$ is *add-bounded* if the non-isomorphic indecomposable direct summands of objects in $\mathcal{C}$ have bounded length. We say that $\mathcal{C}$ is *(n-)syzygy-bounded* if there exists n so that $\Omega^i \mathcal{C}$ is add-bounded for each $i \geq n$. If $A\text{-mod}$ is syzygy-bounded, then we also say that A is syzygy-bounded. Obviously an add-finite (resp., a syzygy-finite) class is add-bounded (resp., syzygy-bounded).

Note that syzygy-bounded classes have finitistic projective dimension by a well-known theorem of Jensen and Lenzing [6] which is based on a theorem of Schofield [9].

It is easy to see that if $\mathcal{D} \subseteq \mathcal{C}$ and $\mathcal{C}$ is add-finite (resp., syzygy-finite, add-bounded, syzygy-bounded), then $\mathcal{D}$ is also add-finite (resp., syzygy-finite, add-bounded, syzygy-bounded).

Now let us recall the definition of Igusa-Todorov algebras introduced in [12]. By definition an artin algebra A is called *(n-)Igusa-Todorov* provided that there exists a fixed A -module ${}_A V$ and a nonnegative integer n such that, for any $M \in A\text{-mod}$, there is an exact sequence $0 \rightarrow V_1 \rightarrow V_0 \rightarrow \Omega_A^n M \rightarrow 0$ with $V_0, V_1 \in \text{add}_A V$. A remarkable property of Igusa-Todorov algebras, which is also the reason why Igusa-Todorov algebras were named, is that these algebras can be shown to satisfy the finitistic dimension conjecture by using the following results on the Igusa-Todorov functor defined in [5].

Lemma 1.1 [5] *For any artin algebra A , there is a functor Ψ which is defined on the objects of $A\text{-mod}$ and takes non-negative integers as values, such that*

- (1) $\Psi(M) = \text{pd}_A M$ provided that $\text{pd}_A M < \infty$.
- (2) $\Psi(X) \leq \Psi(Y)$ whenever $\text{add}_A X \subseteq \text{add}_A Y$. The equation holds in case $\text{add}_A X = \text{add}_A Y$.

(3) If $0 \rightarrow X \rightarrow Y \rightarrow Z \rightarrow 0$ is an exact sequence in $A\text{-mod}$, then $\text{pd}_A Z \leq \Psi(X \oplus Y) + 1$, provided that $\text{pd}_A Z < \infty$.

The class of Igusa-Todorov algebras is large and includes all algebras of representation dimension at most 3, algebras with radical cube zero, monomial algebras and syzygy-finite algebras, etc. [12].

2 Igusa-Todorov and other finiteness conditions

Inspired by the definition of Igusa-Todorov algebras, we introduce here the following notions.

Definition 2.1 Let $\mathcal{C}$ be a class of A -modules. Then $\mathcal{C}$ is said to be $(n\text{-})$ Igusa-Todorov provided that there is an A -module ${}_A V$ and some nonnegative integer n such that, for any $M \in \Omega^n \mathcal{C}$, there is an exact sequence $0 \rightarrow V_1 \rightarrow V_0 \rightarrow M \rightarrow 0$ with $V_0, V_1 \in \text{add}_A V$. The module ${}_A V$ is then called an $(n\text{-})\mathcal{C}$ -Igusa-Todorov module.

By definition, we immediately obtain that all syzygy-finite classes are Igusa-Todorov.

Remark 2.2 (1) If $\mathcal{C} \subseteq \mathcal{D}$ and $\mathcal{D}$ is Igusa-Todorov, then $\mathcal{C}$ is Igusa-Todorov too.

(2) A is an Igusa-Todorov algebra if and only if $A\text{-mod}$ is Igusa-Todorov.

(3) $\mathcal{C}$ is n -Igusa-Todorov if and only if $\Omega^n \mathcal{C}$ is 0-Igusa-Todorov.

(4) If $\Omega^i \mathcal{C}$ is Igusa-Todorov for some $i \geq 1$, then $\mathcal{C}$ is also Igusa-Todorov.

(5) In general, there are many $n\text{-}\mathcal{C}$ -Igusa-Todorov modules for an n -Igusa-Todorov class $\mathcal{C}$.

Lemma 2.3 Let $\mathcal{C}$ be a class of A -modules. If $\mathcal{C}$ is 0-Igusa-Todorov, then $\Omega^i \mathcal{C}$ is also 0-Igusa-Todorov, for any $i \geq 1$.

Proof. Let ${}_A V$ be an $n\text{-}\mathcal{C}$ -Igusa-Todorov module. Then for any $M \in \mathcal{C}$, there is an exact sequence $0 \rightarrow V_1 \rightarrow V_0 \rightarrow M \rightarrow 0$ with $V_0, V_1 \in \text{add}_A V$. By the Horseshoe Lemma, for any $\Omega_A^i M$, there is an exact sequence $0 \rightarrow \Omega_A^i V_1 \rightarrow \Omega_A^i V_0 \oplus P \rightarrow \Omega_A^i M \rightarrow 0$ for some projective A -module P . Hence $\Omega^i \mathcal{C}$ is 0-Igusa-Todorov with a 0- $\Omega^i \mathcal{C}$ -Igusa-Todorov module $\Omega_A^i V \oplus A$. $\square$

Immediately by Lemma 2.3 and Remark 2.2 (3), we obtain the following proposition.

Proposition 2.4 Let $\mathcal{C}$ be a class of A -modules. If $\mathcal{C}$ is n -Igusa-Todorov, then $\Omega^i \mathcal{C}$ is also n -Igusa-Todorov for any $i \geq 1$. In particular $\mathcal{C}$ is m -Igusa-Todorov, for any $m \geq n$ in that case.

The following result gives a method to obtain Igusa-Todorov classes.

Proposition 2.5 Let $\mathcal{C}, \mathcal{D}, \mathcal{E}$ be three classes of A -modules such that, for any $E \in \mathcal{E}$, there is an exact sequence $0 \rightarrow C \rightarrow D \rightarrow E \rightarrow 0$ with $C \in \mathcal{C}$ and $D \in \mathcal{D}$.

(1) If $\mathcal{C}, \mathcal{D}$ are syzygy-finite, then $\mathcal{E}$ is Igusa-Todorov.

(2) If $\mathcal{C}$ is syzygy-finite (resp., syzygy-bounded) and $\text{gd} \mathcal{D} < \infty$, then $\mathcal{E}$ is also syzygy-finite (resp., syzygy-bounded).

(3) If $\mathcal{C}$ is Igusa-Todorov and $\text{gd} \mathcal{D} < \infty$, then $\mathcal{E}$ is also Igusa-Todorov.

Proof. By assumption, for any $E \in \mathcal{E}$, there is an exact sequence $0 \rightarrow C \rightarrow D \rightarrow E \rightarrow 0$ with $C \in \mathcal{C}$ and $D \in \mathcal{D}$. Thus, for any $n \geq 1$, we obtain an exact sequence $0 \rightarrow \Omega^n C \rightarrow \Omega^n D \oplus P \rightarrow \Omega^n E \rightarrow 0$ for some projective A -module P , by the Horseshoe Lemma.

(1) Take n to be an integer such that both $\Omega^n \mathcal{C}$ and $\Omega^n \mathcal{D}$ are add-finite, and denote that $U = \oplus U_i$ (resp., $V = \oplus V_i$), where U_i (resp., V_i) lists all the non-isomorphic indecomposable direct summands of objects in $\Omega^n \mathcal{C}$ (resp., $\Omega^n \mathcal{D}$). Then we have that $\Omega^n C \in \text{add}_A U$ and $\Omega^n D \in \text{add}_A V$. It follows that $\mathcal{E}$ is n -Igusa-Todorov with an n - $\mathcal{E}$ -Igusa-Todorov module $U \oplus V \oplus A$.

(2) Take n to be an integer such that $\Omega^n \mathcal{C}$ is add-finite (resp., add-bounded) and $\text{gd} \mathcal{D} \leq n$. Then $\Omega^n D$ is projective. It follows that $\Omega^n C \simeq \Omega^{n+1} E \oplus Q$ for some projective Q . Now we easily obtain that $\Omega^{n+1} \mathcal{E}$ is add-finite (resp., add-bounded). Hence, $\mathcal{E}$ is syzygy-finite (resp., syzygy-bounded) by the definition involved.

(3) Take n to be an integer such that $\mathcal{C}$ is n -Igusa-Todorov and $\text{gd} \mathcal{D} \leq n$. Similarly to the proof of (2), we obtain that $\Omega^n C \simeq \Omega^{n+1} E \oplus Q$ for some projective Q . Note that there is an exact sequence $0 \rightarrow V_1 \rightarrow V_0 \rightarrow \Omega^n C \rightarrow 0$ with $V_0, V_1 \in \text{add}_A V$, where ${}_A V$ is an n - $\mathcal{C}$ -Igusa-Todorov module. Since Q is projective, we can also obtain an exact sequence $0 \rightarrow V'_1 \rightarrow V'_0 \rightarrow \Omega^{n+1} E \rightarrow 0$ with $V'_0, V'_1 \in \text{add}_A V$. It follows that $\mathcal{E}$ is $(n+1)$ -Igusa-Todorov with ${}_A V$ an $(n+1)$ - $\mathcal{E}$ -Igusa-Todorov module. $\square$

Corollary 2.6 *Let $\mathcal{C}, \mathcal{D}, \mathcal{E}$ be three classes of A -modules such that, for any $E \in \mathcal{E}$, there is an exact sequence $0 \rightarrow C \rightarrow D_0 \rightarrow \cdots \rightarrow D_n \rightarrow E \rightarrow 0$ with $C \in \mathcal{C}$ and each $D_i \in \mathcal{D}$. If $\mathcal{C}$ is syzygy-finite (resp., syzygy-bounded, Igusa-Todorov) and $\text{gd} \mathcal{D} < \infty$, then $\mathcal{E}$ is also syzygy-finite (resp., syzygy-bounded, Igusa-Todorov).*

Proof. Denote $\mathcal{E}_0 := \mathcal{C}_0$, and inductively, $\mathcal{E}_{i+1} := \{M \mid \text{there is an exact sequence } 0 \rightarrow C \rightarrow D \rightarrow M \rightarrow 0 \text{ with } C \in \mathcal{E}_i \text{ and } D \in \mathcal{D}\}$. Then by assumptions and Proposition 2.5, inductively we obtain that each $\mathcal{E}_i$ is syzygy-finite (resp., syzygy-bounded, Igusa-Todorov). Note that $\mathcal{E} \subseteq \mathcal{E}_{n+1}$, so $\mathcal{E}$ is also syzygy-finite (resp., syzygy-bounded, Igusa-Todorov). $\square$

We say that two classes of A -modules $\mathcal{C}, \mathcal{D}$ are *syzygy equivalent* if there exist n, m so that $\text{add}(\Omega^n \mathcal{C}) = \text{add}(\Omega^m \mathcal{D})$ (up to the class $\text{add}_A A$). For example, if there is a finite exact sequence in which all but two modules M, N have finite projective dimension then $\text{add} M, \text{add} N$ are syzygy equivalent. It is obvious that for a class $\mathcal{C}$, all classes $\Omega^i \mathcal{C}$ and $\mathcal{C}$ are syzygy-equivalent. It is also easy to see that $\mathcal{C}$ is syzygy-finite (resp., syzygy-bounded, Igusa-Todorov) if and only if it is in the syzygy equivalent class of an add-finite (resp., add-bounded, 0-Igusa-Todorov) class.

We have the following result.

Proposition 2.7 *Let $A \rightarrow R$ be a homomorphism of artin algebras. If $\text{pd}_A R < \infty$, then the restriction functor:*

$$R - \text{mod} \rightarrow A - \text{mod}$$

preserves syzygy equivalence and sends syzygy equivalence classes of modules to syzygy equivalence classes of modules.

Proof. For a class $\mathcal{C}$ of R -modules, we denote by ${}_A \mathcal{C}$ the image of $\mathcal{C}$ under the restrict functor. Note that for any R -module $M \in \Omega \mathcal{C}$ (resp., $N \in \mathcal{C}$), there is an exact sequence

$0 \rightarrow M \rightarrow P \rightarrow N \rightarrow 0$ with $P \in \text{add}_R R$ and $N \in \mathcal{C}$ (resp., $M \in \Omega\mathcal{C}$). Applying the restrict functor (which is clearly exact), we obtain that, for any A -module ${}_A M \in {}_A\Omega\mathcal{C}$ (resp., ${}_A N \in {}_A\mathcal{C}$), there is an exact sequence of A -modules $0 \rightarrow {}_A M \rightarrow {}_A P \rightarrow {}_A N \rightarrow 0$ with $P \in \text{add}_A R$ and ${}_A N \in {}_A\mathcal{C}$ (resp., ${}_A M \in {}_A\Omega\mathcal{C}$). Since $\text{pd}_A R < \infty$, we see that ${}_A(\Omega\mathcal{C}), {}_A\mathcal{C}$ are syzygy equivalent. It follows that ${}_A(\Omega^i\mathcal{C}), {}_A\mathcal{C}$ are also syzygy equivalent for any $i \geq 0$.

Assume now that $\mathcal{C}, \mathcal{D} \subseteq R\text{-mod}$ are syzygy equivalent. It is enough to show that ${}_A\mathcal{C}, {}_A\mathcal{D} \subseteq A\text{-mod}$ are syzygy equivalent.

By the definition, there exist n, m so that $\text{add}(\Omega^n\mathcal{C}) = \text{add}(\Omega^m\mathcal{D})$ up to the class $\text{add}_R R$. Restricting them to $A\text{-mod}$, we have that $\text{add}_A(\Omega^n\mathcal{C}) = \text{add}_A(\Omega^m\mathcal{D})$ up to the class $\text{add}_A R$. Since $\text{pd}_A R < \infty$, it is easy to see that ${}_A(\Omega^n\mathcal{C}), {}_A(\Omega^m\mathcal{D})$ are syzygy equivalent. Note that ${}_A(\Omega^n\mathcal{C})$ and ${}_A\mathcal{C}$ (resp., ${}_A(\Omega^m\mathcal{D})$ and ${}_A\mathcal{D}$) are syzygy equivalent by the above argument, so we obtain that ${}_A\mathcal{C}, {}_A\mathcal{D}$ are syzygy equivalent. $\square$

Corollary 2.8 *Let $A \rightarrow R$ be a homomorphism of artin algebras such that $\text{pd}_A R < \infty$. If the class of R -modules $\mathcal{C}$ is syzygy-finite (resp., syzygy-bounded, Igusa-Todorov), then its image under the restrict functor, ${}_A\mathcal{C}$, is also syzygy-finite (resp., syzygy-bounded, Igusa-Todorov). In addition, if $\mathcal{D} \subseteq A\text{-mod}$ is syzygy equivalent to a subclass of ${}_A\mathcal{C}$, then $\mathcal{D}$ is also syzygy-finite (resp., syzygy-bounded, Igusa-Todorov).*

Proof. Note that the restrict functor $R\text{-mod} \rightarrow A\text{-mod}$ is exact and length preserving, so all relevant finiteness conditions (e.g., add-finite, add-bounded, 0-Igusa-Todorov) on a class of R -modules are inherited by its image in $A\text{-mod}$. Hence, if $\mathcal{C}$ is syzygy-finite (resp., syzygy-bounded, Igusa-Todorov), then ${}_A\mathcal{C}$ is also syzygy-finite (resp., syzygy-bounded, Igusa-Todorov) by Proposition 2.7. The remained part is obvious. $\square$

Let A and R be artin algebras. Following [14], we say that R is a left idealized extension of A , provided that $A \subseteq R$ has the same identity and $\text{rad}A$ is a left ideal in R . It was shown that every finite dimensional algebra admits a chain of left idealized extensions which is of finite length and ends with a representation-finite algebra [14, Lemma 4.6].

Generalizing [15, Theorem 3.1], it is proved in [12, Theorem 2.9] that if $A = A_0 \subseteq \cdots \subseteq A_m = R$ are artin algebras such that, for each $0 \leq i \leq m-1$, A_{i+1} is a left idealized extension of A_i and $\text{pd}_A A_{i+1} < \infty$, and R is Igusa-Todorov, then the finitistic dimension of A is finite. One part of the following result shows that the artin algebra A obtained in that way is in fact Igusa-Todorov.

Theorem 2.9 *Assume that $A = A_0 \subseteq \cdots \subseteq A_m = R$ are artin algebras such that, for each $0 \leq i \leq m-1$, A_{i+1} is a left idealized extension of A_i and $\text{pd}_A A_{i+1} < \infty$. If R is syzygy-finite (resp., syzygy-bounded, Igusa-Todorov), then A is also syzygy-finite (resp., syzygy-bounded, Igusa-Todorov). In particular, the finitistic dimension of A is finite.*

Proof. Take any $X_0 \in A\text{-mod}$. By [14, Erratum, Lemma 0.1], if A_1 is a left idealized extension of A , then for any $X \in A\text{-mod}$, $\Omega_A^2(X) \simeq \Omega_{A_1} Z \oplus Q$ as A_1 -modules for some $Z, Q \in A_1\text{-mod}$ with ${}_A Q$ projective. Thus, by assumptions and inductively, we see that, for each $0 \leq i \leq m-1$ and each $X_i \in A_i\text{-mod}$, $\Omega_{A_i}^2(X_i) \simeq \Omega_{A_{i+1}}(X_{i+1}) \oplus P_{i+1}$ as A_{i+1} -modules for some $X_{i+1}, P_{i+1} \in A_{i+1}\text{-mod}$ and ${}_A P_{i+1}$ projective. Hence we

have the following exact sequence of A_{i+1} -modules, where P'_{i+1} is the projective cover of $\Omega_{A_{i+1}}(X_{i+1})$:

$$0 \rightarrow \Omega_{A_{i+1}}^2(X_{i+1}) \rightarrow P'_{i+1} \oplus P_{i+1} \rightarrow \Omega_{A_i}^2(X_i) \rightarrow 0.$$

It is easy to see that all these exact sequences can be restricted to the exact sequences of A -modules. Thus we can combine all these exact sequences and obtain the following exact sequence of A -modules:

$$0 \rightarrow \Omega_{A_{m-1}}^2(X_{m-1}) \rightarrow P'_{m-1} \oplus P_{m-1} \rightarrow \cdots \rightarrow P'_1 \oplus P_1 \rightarrow \Omega_A^2(X_0) \rightarrow 0.$$

Denote now $\mathcal{D} = \text{add}_A(\oplus_{i=0}^m A_i)$, then each $P'_i \oplus P_i \in \mathcal{D}$ and $\text{gd}\mathcal{D} < \infty$ by assumption. Let $\mathcal{R}$ be the class of all A -modules which are restricted from R -modules. Then we obtain that $A\text{-mod}$ is syzygy equivalent to a subclass of $\mathcal{R}$ from the above exact sequence. Therefore, if that R is syzygy-finite (resp., syzygy-bounded, Igusa-Todorov), then $A\text{-mod}$ is also syzygy-finite (resp., syzygy-bounded, Igusa-Todorov), by Corollary 2.8. $\square$

Corollary 2.10 *Assume that $A = A_0 \subseteq \cdots \subseteq A_m = R$ are artin algebras such that, for each $0 \leq i \leq m-1$, A_{i+1} is a left idealized extension of A_i and $\text{pd}_{A_i} A_{i+1} < \infty$. If R is syzygy-finite (resp., syzygy-bounded, Igusa-Todorov) and R° has finite finitistic dimension, then the Gorenstein symmetry conjecture, the Wakamatsu-tilting conjecture and the generalized Nakayama conjecture hold for A .*

Proof. Since $\text{pd}_{A_i} A_{i+1} < \infty$ for each $0 \leq i \leq m-1$, we easily obtain that $\text{pd}_A A_{i+1} < \infty$. It follows from Theorem 2.9 that the finitistic dimension of A is finite. Moreover, by assumptions and [13, Corollary 3.7], the finitistic dimension of A° is also finite. Now the conclusion follows from these two facts. $\square$

3 Conditions on ideals

The following result can be compared with [14, Theorem 3.2], where it was proved that if A has two ideals I, J such that $IJ = 0$ and A/I and A/J are representation-finite, then the finitistic dimension of R is finite. In fact, it is shown in [12] that A is 0-Igusa-Todorov in that case.

Theorem 3.1 *Assume that I, J are two ideals of A such that $IJ\text{rad}A = 0$. Then A is Igusa-Todorov provided one of the following conditions is satisfied.*

- (1) A/I is representation-finite, $\text{pd}_A J < \infty$ and A/J is syzygy-finite.
- (2) A/J is representation-finite, $\text{pd}_A I < \infty$ and A/I is syzygy-finite.
- (3) Both A/I and A/J are syzygy-finite and both ${}_A I$ and ${}_A J$ have finite projective dimension.

In particular, the finitistic dimension of A is finite in all three cases.

Proof. Denote by $\mathcal{I}_A$ (resp., $\mathcal{J}_A$) the class of all A -modules which are also A/I -modules (resp., A/J -modules). For any $N \in \Omega(A\text{-mod})$, we have that $N \subseteq \text{rad}_A P$ for some projective ${}_A P$. Denote $C_1 := JN$ and $C_2 := N/JN$, then we have an exact sequence $0 \rightarrow C_1 \rightarrow N \rightarrow C_2 \rightarrow 0$. Note that, by assumptions, $IC_1 = IJN \subseteq IJ\text{rad}_A P = 0$ and

$JC_2 = J(N/JN) = 0$, so C_1 is also an A/I -module and C_2 is also an A/J -module. Now consider the following pullback diagram, where the right column is taken such that ${}_A P$ is the projective cover of ${}_A C_2$.

$$\begin{array}{ccccccc}
& & 0 & & 0 & & \\
& & \downarrow & & \downarrow & & \\
& & \Omega_A C_2 = \Omega_A C_2 & & & & \\
& & \downarrow & & \downarrow & & \\
0 & \rightarrow & C_1 & \rightarrow & C_1 \oplus P & \rightarrow & P \rightarrow 0 \\
& & \parallel & & \downarrow & & \downarrow \\
0 & \rightarrow & C_1 & \rightarrow & N & \rightarrow & C_2 \rightarrow 0 \\
& & & & \downarrow & & \downarrow \\
& & & & 0 & & 0
\end{array}$$

Note that $\Omega_A C_2 \in \Omega \mathcal{J}_A$ and $C_1 \oplus P \in \mathcal{D} := \{C \oplus P \mid C \in \mathcal{I}_A, P \in \text{add}_A A\}$, so if we show that both $\Omega \mathcal{J}_A$ and $\mathcal{D}$ are syzygy-finite, then A is Igusa-Todorov by Remark 2.2 (3), Proposition 2.5 and the middle column in the diagram.

(1) If $\text{pd}_A J < \infty$ and A/J is syzygy-finite, then $\text{pd}_A A/J < \infty$. By Corollary 2.8, we obtain that the class $\mathcal{J}_A$ is syzygy-finite. It follows that $\Omega \mathcal{J}_A$ is also syzygy-finite. Since A/I is representation-finite, the class $\mathcal{D}$ is clearly also syzygy-finite.

(2) Similarly as in (1), we have that $\mathcal{I}_A$ is syzygy-finite. It follows that $\mathcal{D}$ is also syzygy-finite. Since A/J is representation-finite, the class $\Omega \mathcal{J}_A$ is clearly also syzygy-finite.

(3) Similarly as in (1), we have that both $\mathcal{I}_A$ and $\mathcal{J}_A$ are syzygy-finite by Corollary 2.8. Consequently, both $\Omega \mathcal{J}_A$ and $\mathcal{D}$ are syzygy-finite. $\square$

The following list mentions some known classes of syzygy-finite algebras.

- Algebras of finite representation type (in fact 0-syzygy-finite).
- Algebras of finite global dimension.
- Monomial algebras (indeed 2-syzygy-finite) [17].
- Left serial algebras (indeed 2-syzygy-finite) [18].
- Torsionless-finite algebras (indeed 1-syzygy-finite) c.f. [8], including:
 - Algebras A with $\text{rad}^n A = 0$ and $A/\text{rad}^{n-1} A$ representation-finite.
 - Algebras A with $\text{rad}^2 A = 0$.
 - Minimal representation-infinite algebras.
 - Algebras stably equivalent to hereditary algebras.
 - Right glued algebras and left glued algebras.
 - Algebras of the form $A/\text{soc} A$ with A a local algebra of quaternion type.
 - Special biserial algebras.
- Algebras possessing a left idealized extension which is torsionless-finite (indeed 2-syzygy-finite) [12].

Corollary 3.2 *Let A be an artin algebra. If there is an ideal I such that $I^2 \text{rad} A = 0$ (for instance $I^2 = 0$), $\text{pd}_A I < \infty$ and A/I is syzygy-finite, then the finitistic dimension of A is finite.*

A special case of the above corollary is as follows.

Corollary 3.3 *Let A be an artin algebra with $\text{rad}^{2n+1}A = 0$. If $\text{pd}_A \text{rad}^n A < \infty$ and $A/\text{rad}^n A$ is syzygy-finite, then the finitistic dimension of A is finite.*

In particular, we obtain the following result.

Corollary 3.4 *Let A be an artin algebra with $\text{rad}^5 A = 0$. If $\text{pd}_A \text{rad}^2 A < \infty$, then the finitistic dimension of A is finite.*

Proof. Note that $A/\text{rad}^2 A$ is an algebra with radical square zero and hence is torsionless-finite. Thus the conclusion follows from Corollary 3.3. $\square$

Proposition 3.5 *Let A be an artin algebra and I be an ideal with $I\text{rad}A = 0$ and $\text{pd}_A I < \infty$. If A/I is syzygy-finite (resp., syzygy-bounded, Igusa-Todorov), then A is also syzygy-finite (resp., syzygy-bounded, Igusa-Todorov). In particular, the finitistic dimension of A is finite.*

Proof. By assumptions, for any $N \in A\text{-mod}$, we have that $I\Omega_A N \subseteq I\text{rad}(P_N) = 0$, where P_N is the projective cover of N . Hence $\Omega_A N \in \mathcal{I}_A$, where $\mathcal{I}_A$ denotes the class of all A -modules which are also A/I -modules. It follows that $A\text{-mod}$ is syzygy equivalent to a subclass of $\mathcal{I}_A$. Moreover, we have that $\text{pd}_A A/I < \infty$, since $\text{pd}_A I < \infty$ by the assumption. Hence, if A/I is syzygy-finite (resp., syzygy-bounded, Igusa-Todorov), then $A\text{-mod}$ is also syzygy-finite (resp., syzygy-bounded, Igusa-Todorov) by Corollary 2.8. $\square$

Generally, we have the following result.

Theorem 3.6 *Assume that $0 = I_0 \subseteq I_1 \subseteq \cdots \subseteq I_n$ are ideals in A such that $\text{pd}_A I_{i+1} < \infty$ and $(I_{i+1}/I_i)\text{rad}(A/I_i) = 0$, for each $0 \leq i \leq n-1$. If A/I_n is syzygy-finite (resp., syzygy-bounded, Igusa-Todorov), then A is also syzygy-finite (resp., syzygy-bounded, Igusa-Todorov). In particular, the finitistic dimension of A is finite.*

Proof. Denote that $A_i = A/I_i$ for each $0 \leq i \leq n$. So $A_0 = A$. Note that $\text{pd}_A A_i < \infty$ for each $0 \leq i \leq n$, since $\text{pd}_A I_i < \infty$ by assumption.

Take any $X_0 \in A_0\text{-mod}$ and denote $X_1 = \Omega_{A_0} X_0$, then we have an exact sequence $0 \rightarrow X_1 \rightarrow P_{X_0} \rightarrow X_0 \rightarrow 0$, where P_{X_0} is the projective cover of X_0 . Since $I_1 X_1 = I_1 \Omega_{A_0} X_0 \subseteq I_1 \text{rad}_A(P_{X_0}) = 0$ by assumptions, we see that X_1 is also an A_1 -module. Inductively, for each $0 \leq i \leq n-1$ and each A_i -module X_i , we obtain an exact sequence $0 \rightarrow X_{i+1} \rightarrow P_{X_i} \rightarrow X_i \rightarrow 0$, where P_{X_i} is the projective cover of the A_i -module X_i , such that X_{i+1} is also an A_{i+1} -module.

Now restricting these exact sequences to A -modules and combining them, we obtain an exact sequence $0 \rightarrow X_n \rightarrow P_{X_{n-1}} \rightarrow \cdots \rightarrow P_{X_0} \rightarrow X_0 \rightarrow 0$ of A -modules, where X_n is also an A_n -module and each P_{X_i} is projective as an A_i -module.

Denote now $\mathcal{D} = \text{add}_A(\bigoplus_{i=0}^{n-1} A_i)$, then each $P_{X_i} \in \mathcal{D}$ and $\text{gd}\mathcal{D} < \infty$ by assumption. Let $\mathcal{A}_n$ be the class of all A -modules which are also A_n -modules. Then we obtain that $A\text{-mod}$ is syzygy equivalent to a subclass of $\mathcal{A}_n$ from the above exact sequence. Now the conclusion easily follows from Corollary 2.8. $\square$

Corollary 3.7 *Let n be the integer such that $\text{rad}^n A = 0$.*

(1) *If $\text{pd}_A \text{rad}^i A < \infty$ for each $i \geq 3$, then A is Igusa-Todorov.*

(2) *If $\text{pd}_A \text{rad}^i A < \infty$ for each $i \geq 2$, then A is syzygy-finite.*

In particular, the finitistic dimension of A is finite in either case.

Proof. Consider the ideals $0 = \text{rad}^n A \subseteq \text{rad}^{n-1} A \subseteq \cdots \subseteq \text{rad}^3 A \subseteq \text{rad}^2 A$. Then clearly $(\text{rad}^{i-1} A / \text{rad}^i A) \cdot \text{rad}(A / \text{rad}^i A) = 0$.

(1) Since $A / \text{rad}^3 A$ is an algebra with radical cube zero, it is Igusa-Todorov by [12]. Hence we obtain the conclusion by assumptions and Theorem 3.6.

(2) Since $A / \text{rad}^2 A$ is an algebra with radical square zero, it is syzygy-finite. Again we obtain the conclusion by assumptions and Theorem 3.6. $\square$


As special cases of Corollaries 3.4 and 3.7, we obtain the following result.

Corollary 3.8 *If $\text{rad}^4 A = 0$, then the finitistic dimension of A is finite provided that $\text{pd}_A \text{rad}^2 A < \infty$ or $\text{pd}_A \text{rad}^3 A < \infty$.*

4 Examples

In this section, we will present examples to show that our results do apply to check the finiteness of the finitistic dimensions of some algebras.

Example 4.1 [4] *Consider the bound quiver algebra $A = kQ/J$ where Q is given by*




and J is generated by the set of paths $\{\alpha^3, \beta\alpha, \alpha\mu_i\rho, \beta\mu_i, \delta\gamma_1 - \delta\gamma_2\}$ where $i = 1, 2$. Then A is syzygy-finite. In particular, the finitistic dimension of A is finite.

The algebra was proved to have finite finitistic dimension in [4] by showing that $\ell\ell^\infty A = 3$. Here the notion $\ell\ell^\infty A$ denotes the infinite-layer length of A , see [4] for details. In fact, one main result in [4] asserts that an artin algebra A has finite finitistic dimension whenever $\ell\ell^\infty A \leq 3$, generalizing the known result for algebras with radical cube zero.

Now we give a different proof of the finiteness of the finitistic dimension of A . Moreover, we prove that A is syzygy-finite.

To see this, let I_1 be the ideal generated by β and I_2 be the ideal generated by $\beta, \gamma_1, \gamma_2, e_2$. Then $I_1 \subseteq I_2$ and $I_1 \text{rad} A = 0 = (I_2 / I_1) \text{rad}(A / I_1)$. Moreover, $\text{pd}_A I_1 < \infty$ and $\text{pd}_A I_2 < \infty$. Note that A / I_2 is of the form




with relations $0 = \alpha^3 = \alpha\mu_i\rho$, where $i = 1, 2$. Clearly A/I_2 is a monomial algebra and hence is syzygy-finite. Now by Theorem 3.6 we obtain that A is also syzygy-finite.

Example 4.2 Consider the bound quiver algebra $A = kQ/J$ where Q is the same as in Example 4.1, but J is generated by the set of paths $\{\alpha^6, \beta\alpha, \alpha\mu_i\rho, \beta\mu_i, \delta\gamma_1 - \delta\gamma_2\}$, where $i = 1, 2$. Then A is syzygy-finite. In particular, the finitistic dimension of A is finite.

The algebra is revisited from Example 4.1 and the argument is the same. Note that here $\ell\ell^\infty A = 6$, so we can't use the results in [4].

Example 4.3 [16] Consider the bound quiver algebra $A = kQ/J$ where Q is given by




and J is generated by the set of paths $\{\alpha\beta, \alpha\epsilon, \alpha\delta, \alpha\xi, \gamma\beta, \gamma\delta, \gamma\epsilon, \gamma\xi, \zeta\epsilon, \epsilon\zeta, \xi\zeta, \eta\epsilon, \eta\xi, \beta\alpha - \delta\gamma\}$. Then A is syzygy-finite. In particular, the finitistic dimension of A is finite.

It was shown in [16] that A has a semisimple extension A' such that $\text{rad}A = \text{rad}A'$ and A' is a trivially twisted extension of two algebras of finite finitistic dimension. Consequently, by results in [16], the algebra A has finite finitistic dimension, see [16] for details.


Now we give a different proof of the finiteness of the finitistic dimension of A . Moreover, we also show that A is syzygy-finite.

To see this, let I be the ideal generated by α . Then $I\text{rad}A = 0$. Moreover, $\text{pd}_A I < \infty$. Note that A/I is of the form



with relations $0 = \delta\gamma = \gamma\beta = \gamma\delta = \gamma\epsilon = \gamma\xi = \zeta\epsilon = \epsilon\zeta = \xi\zeta = \eta\epsilon = \eta\xi$. Clearly A/I is a monomial algebra and hence is syzygy-finite. Now by Proposition 3.5 we obtain that A is also syzygy-finite.


Example 4.4 Consider the bound quiver algebra $A = kQ/J$ where Q is given by



and J is generated by paths $\{\rho^6, \sigma^6, \alpha\beta, \alpha\epsilon, \alpha\delta, \alpha\xi, \gamma\beta, \gamma\delta, \gamma\epsilon, \gamma\xi, \zeta\epsilon, \epsilon\zeta, \xi\zeta, \eta\epsilon, \eta\xi, \beta\alpha - \delta\gamma\}$. Then A is syzygy-finite. In particular, the finitistic dimension of A is finite.


The algebra is revisited from Example 4.3 and the argument is almost the same.

Example 4.5 Consider the bound quiver algebra $A = kQ/J$ where Q is given by



and J is generated by the set of paths $\{\alpha\gamma_i\beta - \alpha\gamma_j\beta \text{ for all } 1 \leq i < j \leq n, \alpha^3, \beta^6\}$. Then A is Igusa-Todorov. In particular, the finitistic dimension of A is finite.

In fact, let I be the ideal generated by $\{\gamma_i, 1 \leq i \leq n-1\}$. Then $I^2 = 0$. Moreover, $\text{pd}_A I < \infty$. Note that A/I is of the form



with relations $0 = \alpha\gamma\beta = \alpha^3 = \beta^6$. Clearly A/I is a monomial algebra and hence is syzygy-finite. Now by Corollary 3.2 we obtain that A is Igusa-Todorov.

Similarly, we can also obtain that A° is also Igusa-Todorov and hence has also finite finitistic dimension.

We note that the Gorenstein symmetry conjecture, the Wakamatsu-tilting conjecture and the generalized Nakayama conjecture also hold for the algebras in these examples.

ACKNOWLEDGEMENTS

The author thanks for the referee's highly valuable comments and suggestions. In particular, the notions of "syzygy-bounded" and "syzygy-equivalent" are due to the referee, as well as Propositions 2.7.

References

- [1] Green E. L., Kirmann E. and Kuzmanovich J., Finitistic dimensions of finite monomial algebras, J. Algebra 136 (1991), 37-50.

- [2] Green E. L. and Zimmermann-huisgen B., Finitistic dimension of artinian rings with vanishing radical cube, *Math. Z.* 206 (1991), 505-526.
- [3] Happel D., Homological conjectures in representation theory of finite- dimensional algebras, *Sherbrook Lecture Notes Series* (1991), available from <http://www.math.ntnu.no/~oyvinso/Nordfjordeid/Program/references.html>.
- [4] Huard F., Lanzilotta M. and Mendoza O., Finitistic dimension through infinite projective dimension, *Bull. London Math. Soc.* 41 (2009), 367-376.
- [5] Igusa K. and Todorov G., On the finitistic global dimension conjecture for artin algebras. In: *Representations of algebras and related topics*, 201-204. *Fields Inst. Commun.* 45, Amer. Math. Soc., Providence, RI, 2005.
- [6] Jensen C. U. and Lenzing H., *Model theoretic algebra*. New York 1989.
- [7] Mantese F. and Reiten I., Wakamatsu Tilting modules, *J. Algebra* 278 (2004), 532-552.
- [8] Ringel C.M., The torsionless modules of an artin algebra, preprint (2008).
- [9] Schofield A. H., Bounding the global dimension in terms of the dimension, *Bull. London Math. Soc.* 17. 393-394 (1985).
- [10] Small L. W., A change of rings theorem. *Proc. Amer. Math. Soc.* 19(1968), 662-666.
- [11] Wei J., A note on Auslander bounds, [arXiv:math.RT/0802.1085](https://arxiv.org/abs/math.RT/0802.1085).
- [12] Wei J., Finitistic dimension and Igusa-Todorov algebras, preprint.
- [13] Wei J., Finitistic dimension and restricted flat dimension, *J. Algebra* 320 (2008), 116-127.
- [14] Xi C., On the finitistic dimension conjecture I: related to representation-finite algebras, *J. Pure and Appl. Algebra* 193 (2004), 287-305. Erratum: *J. Pure and Appl. Algebra* 202 (1-3) (2005), 325-328.
- [15] Xi C., On the finitistic dimension conjecture II: related to finite global dimension, *Adv. Math.* 201 (2006), 116-142.
- [16] Xi C. and Xu D., On the finitistic dimension conjecture IV: related to relatively projective modules, preprint.
- [17] Zimmermann-Huisgen B., Homological domino effects and the first finistic demension conjecture, *Invent. Math.* 108 (1992), 369-383.
- [18] Zimmermann-Huisgen B., Syzygies and homological dimensions over left serial rings, in *Methods in Module Theory* (G. Abrams, J. Haefner, and K.M. Rangaswamy, Eds.), pp. 161-174, New York (1992) Dekker.
- [19] Zimmermann-Huisgen B., The finitistic dimension conjectures - A tale of 3.5 decades, In: *Abelian Groups and Modules*, 501-517, Kluwer, Dordrecht, 1995.